

**A CARLAT PSYCHIATRY  
REFERENCE TABLE**

<b>Intravenous Microinduction Strategies Utilized</b>		
<b>Dosing Strategy</b>	<b>Hour of Treatment</b>	<b>Buprenorphine Dose</b>
<b>Rapid</b>	0-6	0.3 mg IV q6h X 2 doses
	12-18	0.6 mg IV q6h X 2 doses
	24	4 mg SL X 1 dose
<b>Moderate</b>	0-6	0.15 mg IV q6h X 2 doses
	12-18	0.3 mg IV q6h X 2 doses
	24-30	0.6 mg IV q6h X 2 doses
	36	4 mg SL X 1 dose
<b>Slow</b>	0-18	0.15 mg IV q6h X 4 doses
	24-30	0.3 mg IV q6h X 2 doses
	36-42	0.6 mg IV q6h X 2 doses
	48	4 mg SL X 1 dose

Source: Jablonski LA et al, Drug Alcohol Depend 2022;237:109541

From the Research Update:  
**“Intravenous Buprenorphine Microinduction”**  
 By Marissa Bayerl, APRN, and Noah Capurso, MD, MHS, Ms  
*The Carlat Addiction Treatment Report*, Volume 11, Number 5&6 July/August/September  
[www.carlataddictiontreatment.com](http://www.carlataddictiontreatment.com)