A CARLAT PSYCHIATRY REFERENCE TABLE

Symptoms of Infective Endocarditis	
Common	Rare, but Specific
Fever New-onset cardiac murmur Nonspecific constitutional symptoms	 Janeway lesions: Painless red spots on palms and soles of feet Osler nodes: Raised tender lesions on palms and soles of feet Roth spots: Retinal hemorrhages with white centers Splinter hemorrhages: Vertical areas of bleeding under nail beds
Medical Complications, Late Disease	
 Acute kidney injury: Decreased urinary output, edema, confusion, fatigue Congestive heart failure: Shortness of breath, exercise intolerance, edema Pulmonary embolus/infarct: Shortness of breath, chest pain, hemoptysis Sepsis: High or low body temperature, tachycardia, tachypnea, leukocytosis Splenic infarct: Left-sided abdominal pain, splenomegaly, nausea, vomiting Stroke: Weakness, speech difficulty, facial palsy, loss of balance 	

From the Clinical Update:
"What Psychiatrists Need to Know About Intravenous Drug Use and Endocarditis"
By Kenechukwu Mezue, MD, MSc, and Bernardo Lombo, MD, FACC
The Carlat Addiction Treatment Report, Volume 11, Number 5&6 July/August/September
www.carlataddictiontreatment.com