A CARLAT PSYCHIATRY REFERENCE TABLE

Comparison of Brixadi vs Sublocade		
Feature	Brixadi	Sublocade
Dosing Options	• 7 dose options (weekly & monthly)	• 2 dose options (monthly only)
• Formulations	Weekly and monthly	• Monthly only (can be given every other month)
Induction Timing	Can be started earlier in withdrawal, useful in acute settings	Requires moderate with- drawal symptoms and at least one sublingual dose before initiation
Injection Volume	• Smaller (0.16–0.64 mL); less painful	• Larger (0.5–1.5 mL)
Needle Size	• Thinner 23-gauge needle	Thicker 19-gauge needle
Injection Experience	• Faster injection (a few seconds); less painful	• Slower injection (30–60 seconds); more discomfort
Patient Preference	Higher satisfaction in some studies; less injection site pain	• Lower preference among patients with experience of both
Flexibility for Monitoring	Weekly dosing allows closer follow-up for high-need patients	Monthly/every-other-month better for stable patients
• Induction in Acute Settings	Suitable due to low risk of precipitated withdrawal	• Less ideal for rapid induction

From the Clinical Update:

"Brixadi: A New Long-Acting Buprenorphine Option for Opioid Use Disorder"

By Crystal Obiozor, MD and Noah Capurso, MD, MHS
The Carlat Addiction Treatment Report, Volume 13, Number 5&6, July/August/September 2025

www.carlataddictiontreatment.com