

A CARLAT PSYCHIATRY REFERENCE TABLE

Medications for Stimulant Use Disorders		
Medication	Dosing	Special Considerations
Bupropion XL	450 mg PO qAM	<ul style="list-style-type: none"> • Good for comorbid ADHD, depression, nicotine use disorder • Risk of seizures • Risk of abuse • Combine with naltrexone (XR IM q3wks)
Disulfiram	250–500 mg PO daily	<ul style="list-style-type: none"> • Good for comorbid AUD • Risk of hepatotoxicity • Beware inadvertent reaction with alcohol, especially at high doses
Mirtazapine	30 mg PO daily	<ul style="list-style-type: none"> • Good for comorbid depression, insomnia, underweight
Modafinil	400 mg PO qAM	<ul style="list-style-type: none"> • Good for comorbid ADHD • Risk of abuse
Naltrexone	380 mg IM q3wks	<ul style="list-style-type: none"> • Good for comorbid AUD • Caution in patients with liver disease • Combine with bupropion
Topiramate	200–300 mg PO daily	<ul style="list-style-type: none"> • Good for comorbid AUD and weight loss • Cognitive clouding side effect • Risk of kidney stones, metabolic derangements

From the Clinical Update:
“Pharmacotherapy for Stimulant Use Disorders”

By **Rehan Aziz, MD**

The Carlat Addiction Treatment Report, Volume 9, Number 3, May/June 2021

www.carlataddictiontreatment.com