Agents Used in Medication-Assisted Treatment (MAT) for Opioid Addiction Generic Name (Brand Name) Formulation and Usual Advantages/Disadvantages Year FDA Approved (Rx status) **Available Strengths** Dosage Range (mg) [G] denotes generic availability Buprenorphine Sublingual tablets: 2, 8 4-24 QD Advantages: Original mono version; has a long track (formerly Subutex) [G] record; very inexpensive. Many experts find it to be just as effective as agents that include naloxone. 2002 (C-III) Disadvantage: Since it does not contain naloxone, can be ground up and injected. Buprenorphine and naloxone [G] Sublingual tablets: 4-24 OD Advantages: First combination buprenorphine/nalox-Bup 2/Nx 0.5, 8/2 one agent. Long track record; inexpensive. 2002 (C-III) Disadvantage: Some complain about the taste. Buprenorphine and naloxone Sublingual strips: Advantages: Faster absorption than tablets; easy to 4-24 OD (Suboxone) [G] Bup 2/Nx 0.5, taper gradually because film can be cut into small 4/1, 8/2, 12/3 sizes; packaging makes it more difficult for kids to 2010 open. (C-III) Disadvantages: More diversion potential because you can easily mail the strips; relatively high cost. Buprenorphine and naloxone 2.1-12.6 QD Advantages: High bioavailability, fast absorption, less Buccal film: (Bunavail) Bup 2.1/Nx 0.3, constipation; more convenient than other preparations 4.2/0.7, 6.3/1 because it sticks to the cheek while dissolving, allow-2014 (C-III) ing patients to talk. Disadvantages: Very high cost; may be more difficult to cut than Suboxone film, because cutting may decrease its sticking ability to cheek mucosa. Buprenorphine and naloxone Sublingual tablets: 2.9-17.2 QD Advantages: Menthol flavor; higher bioavailability than (Zubsolv) Bup 1.4/Nx 0.36, generics. 2013 2.9/0.71, 5.7/1.4, Disadvantage: Higher cost than generics. (C-III) 8.6/2.1, 11.4/2.9 Methadone [G] Oral tablets: 5, 10, 40 20-120 OD Advantages: Long track record; daily dosing require-Oral liquid: 10 mg/mL, ment may be good for those needing closer supervi-(Dolophine, Methadose) 10 mg/5 mL, 5 mg/5 mL 1947 sion. (C-II) Disadvantages: Daily dosing is inconvenient for those with job or family duties; many drug interactions; high risk of respiratory depression. 380 Q 4wk Advantages: Monthly injection decreases worries Naltrexone ER IM injection: 380 (Vivitrol) about compliance; beneficial for alcohol use disorder. 2006 Disadvantages: Shot can be painful; patients may not (Rx) return for repeat injections; patients must go through longer period of withdrawal before starting treatment.

Buprenorphine/naloxone products are not bioequivalent. Comparable doses: 2mg/0.5 mg SL strip = 1.4 mg/0.36 mg SL tab; 4 mg/1 mg SL strip = 2.1 mg/0.3 mg buccal film; 8 mg/2mg SL strip = 4.2 mg/0.7 mg buccal film or 5.7 mg/1.4 mg SL tab; 12 mg/3 mg SL strip = 6.3 mg/1 mg buccal film